



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 1 013 752 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
28.06.2000 Bulletin 2000/26

(51) Int. Cl.<sup>7</sup>: **C11B 1/00, A23D 9/00,  
A23D 7/00, A23L 1/24**

(21) Application number: **99204127.7**

(22) Date of filing: **06.12.1999**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: **23.12.1998 EP 98204443**

(71) Applicants:  
• **UNILEVER N.V.**  
**3013 AL Rotterdam (NL)**  
• **UNILEVER PLC**  
**London EC4P 4BQ (GB)**

(72) Inventor:  
**van Putte, Karel Petrus A. M.,  
Unilever Res VI.  
3133 AT Vlaardingen (NL)**

(74) Representative:  
**Sikken, Antonius H. J. M. et al  
UNILEVER N.V.,  
Patent Division,  
P.O. Box 137  
3130 AC Vlaardingen (NL)**

(54) **Process for preparing food products fortified with oleanolic acid**

(57) Olive oil is fortified with oleanolic acid by including olive leaves in the malaxation mash of conventional olive oil pressing in an amount of 5 - 50 wt.%, based on the weight of the malaxation mash.

The invention comprises vegetable oils fortified with a concentrate of oleanolic acid and, optionally, oleuropein. Food products fortified with oleanolic acid containing oils or concentrates according to the invention are also part of the invention.

**EP 1 013 752 A1**

## Description

[0001] The present invention relates to a process for preparing food products containing a high level of oleanolic acid, a compound which is derived from leaves of the olive tree. The present invention also relates to the obtained fortified products such as vegetable oils, in particular olive oil, spreads, mayonnaise, salad dressings and sauces.

## BACKGROUND OF THE INVENTION

[0002] Oleanolic acid (3-beta-hydroxy-28-carboxy-oleanen) belongs to the chemical compound class of the triterpenes. See Figure 1 for its chemical structure. Olive leaves are a known source of oleanolic acid, containing up to 6.5 wt.%. Oleanolic acid has a low solubility in olive oil and may cause turbidity.

[0003] Oleanolic acid has a low solubility in olive oil and may cause turbidity.

[0004] Japanese Patent JP 9078061 (Derwent Abstract) discloses the use of a hydrolysed extract of olives and olive leaves as an antioxidant to be used for food, feed, cosmetics and drugs.

[0005] J. of the Am. Oil Chemists Soc. Vol. 73, no.3, pp. 371-374 (1996) gives a description of harvesting olive oil by pressing olive fruits at ambient temperature. For improving the olive oil flavour some olive leaves are added to the process mixture.

[0006] It is known that olive oil contains low amounts of oleanolic acid, although its presence is avoided because its low solubility in oil causes turbidity, particularly when the oil is stored below ambient temperature.

[0007] Another useful component of olive oil is oleuropein, one of the native phenolic compounds which because of its bitter taste is removed from olive oil after pressing by washing or hydrolysis. Presently it is appreciated for its health supporting properties. Oleuropein is believed to have anti-inflammatory and anti-oxidant properties.

[0007] In olive oil which is obtained by conventional olive oil processing no levels higher than 300 ppm for oleanolic acid have been reported. Generally, olive oil is prepared by harvesting olive fruits and subjecting these to a malaxation treatment which comprises crushing and kneading the olives so that a mash is obtained containing the whole content of the olive fruit: solid matter, oil and an aqueous phase. Accidentally, some leaves of the olive tree may be encountered in the malaxation mixture. After malaxation the mash is separated into a liquid phase and a solid residue. The olive oil is separated from the aqueous phase in a decanting vessel or by centrifugation. The olive oil may be washed with water.

## SUMMARY OF THE INVENTION

[0008] We have discovered the fortification of food products with oleanolic acid. The nutritional potential of this compound is based on its many valuable health supporting properties (see EP 94 517). The invention provides an economical and simple process for the fortification of food products with oleanolic acid, in particular of a vegetable oil such as olive oil. According to the present invention an olive oil or olive residue oil is prepared according to a conventional process, but with the exception that

- the malaxation mash contains 5 - 50 wt.% of olive leaves
- the malaxation treatment is carried out at an elevated temperature being 24-60°C.

This adapted process results in an olive oil or an olive residue oil which contains at least 300 ppm of oleanolic acid.

## DETAILS OF THE INVENTION

[0009] The present invention relates to a process for preparing an olive oil fortified with oleanolic acid, which comprises the steps of:

- subjecting a mixture of olive fruits and olive leaves to a malaxation treatment resulting in a mash which contains a first olive oil and a first residue; and
- separating the first olive oil from the first residue.

[0010] In order to get a substantial amount of oleanolic acid dissolved, the malaxation process has to be carried out at an elevated temperature being 24-60°C, preferably 30-60°C, more preferably 30-40°C.

In the malaxation mixture olive leaves are present in an amount 5 - 50 wt.%, preferably 10 - 50 wt.%, based on the weight of the mash.

[0011] Since the leaves of the olive are such a rich source of oleanolic acid, the present process results in a substantial fortification of the olive oil. The extent of fortification not only depends on the amount of added leaves and extraction temperature, but also on the type of the olive tree. The content of oleanolic acid in the leaves may vary from

year to year. For obtaining a specific content of oleanolic acid in the olive oil, a suitable approach is first to determine the amount of this compound in the olive leaves and then to adjust the amount of olive leaves added to the malaxation mixture.

[0012] The process according to the invention may include conventional process steps, such as water washing of olives and leaves. Water may be added before the first olive oil is separated from the first residue.

Decantation, centrifugation and filtration are suitable processes for the separation of olive oil from residue and aqueous phase.

[0013] According to a further embodiment of the present invention water is added to the first residue and the residue is extracted employing conventional extraction conditions. This process yields a second olive oil and a second residue. The second olive oil is separated from the second residue. The second olive oil also contains a considerable amount of oleanolic acid. Preferably, before extraction extra olive leaves are added to the first residue, preferably, in an amount of 5 - 50 wt.%, more preferably 10 - 50 wt.%, based on the total extraction mass.

[0014] According to still another embodiment of the invention the olive leaves are not added to the fresh olives but only to the first residue.

[0015] Subsequently, one or both residues can be subjected to solvent extraction, applying conventional process conditions, which yields an olive residue oil which is also fortified with oleanolic acid. From the separated mixture of solvent and oil the fortified olive residue oil is obtained by evaporating the solvent. Suitable solvents are hexane and acetone.

TABLE I

	Oleanolic acid (ppm)
Olive oil obtained by conventional process	< 300
Olive oil obtained with malaxation at 30°C in the presence of 50 wt.% of olive leaves	400

[0016] According to another embodiment of the invention an oleanolic acid concentrate is prepared starting with the olive oil or olive residue oil obtained with the present invention.

The fortified olive oil or olive residue oil is subjected to an extraction with an oil immiscible solvent which results into a solvent phase containing oleanolic acid and an oil phase. By distilling the solvent from the solvent phase the desired concentrate is obtained. Suitable solvents for oleanolic acid are alcohols, preferably methanol and ethanol and mixtures of an alcohol with water.

[0017] The present invention comprises an olive oil or an olive residue oil which contains at least 300 ppm, preferably at least 350 ppm, more preferably at least 450 ppm of oleanolic acid.

[0018] The invention further comprises an olive oil or an olive residue oil which contains at least 300 ppm, preferably at least 350 ppm, more preferably at least 450 ppm of oleanolic acid and additionally at least 20 ppm, preferably at least 100 ppm of oleuropein.

The oleuropein may be a native component of the oil or it may be added in isolated form.

[0019] The effect of high concentrations of oleanolic acid in the obtained oil is the occurrence of turbidity when after processing the oil cools down to ambient temperature. The turbidity is caused by dispersed, not dissolved oleanolic acid. Although turbidity is not acceptable when a clear olive oil is desired, it does not harm when the oil is used as an ingredient for preparing a food product according to the invention.

[0020] The olive oil, olive residue oil and concentrate obtained according to the present invention are useful ingredients for the preparation of fortified food products, preferably olive oils, such as extra virgin olive oil, fine, semi-fine or regular virgin olive oil, refined olive oil, or olive residue oil and blends which contain virgin olive oil and refined olive oil.

[0021] The oleanolic acid concentrate can be admixed e.g. with other vegetable oils such as rapeseed oil, sunflowerseed oil, soybean oil and corn oil.

Common food products such as spreads, mayonnaises, salad dressings and sauces can be fortified by substituting the oil component by an oil according to the invention. Spreads are understood to be products which contain a substantial amount of fat, often 40 wt.% and more, which have a plastic consistency and which are used for buttering bread.

## Claims

1. A process for fortifying olive oil with oleanolic acid, which comprises the steps of:

- subjecting harvested olives to a malaxation treatment,
- separating the malaxation mash into an olive oil phase denoted as first olive oil, a solid residue denoted as first

- residue and an aqueous phase,
- collecting the olive oil,

characterized in that

the malaxation mash contains, calculated on mash, 5 - 50 wt.% of olive leaves and that the malaxation treatment is carried out at 24-60°C.

2. A process according to claim 1, characterized in that the malaxation is carried out at 30-60°C, preferably at 30-40°C.

3. A process according to claims 1 or 2, characterized in that the mash contains 5 - 50 wt.%, preferably 10 - 50 wt.% of olive leaves.

4. A process according to anyone of claims 1 - 3, which comprises the further step of:

extracting the first residue with water resulting in a mixture of a second olive oil and a second residue, separating the second olive oil from the second residue and collecting the second olive oil.

5. A process according to claim 4, wherein the first residue before being extracted with water is mixed with, calculated on total extraction mass, 5 - 50 wt.%, preferably with 10 - 50 wt.% of olive leaves.

6. A process for fortifying an olive oil with oleanolic acid, which comprises the steps of:

- subjecting a mash of crushed olives which contains olive leaves to a malaxation treatment,
- separating the mash after malaxation into an olive oil phase denoted as first olive oil, a solid residue denoted as first residue and an aqueous phase,
- adding olive leaves to the first residue in an amount of 5 - 50 wt.%, preferably 10 - 50 wt.%, based on the weight of the extraction mass,
- extracting with water the first residue containing the leaves to obtain a mixture of a second olive oil and a second residue and
- separating the second olive oil from the second residue.

7. A process for preparing an olive residue oil fortified with oleanolic acid, which comprises the steps of:

- subjecting a residue obtained with a process according to any of the claims 1 - 6 to solvent extraction to obtain a mixture of oil, solvent and a final residue,
- separating the liquid phase from the solid residue phase - removing the solvent by evaporation from the oil to obtain an olive residue oil.

8. A process for preparing a concentrate containing oleanolic acid, which comprises the steps of:

- subjecting the oil obtained with a process according to any of the claims 1 - 7, to an extraction with an oil immiscible solvent in which oleanolic acid can dissolve to obtain a solvent phase which contains oleanolic acid,
- evaporating the solvent from the solvent phase and
- collecting a concentrate containing oleanolic acid.

9. Olive oil which contains at least 300 ppm, preferably at least 350 ppm, more preferably at least 450 ppm of oleanolic acid.

10. Olive oil according to claim 9 which additionally contains at least 20 ppm, preferably at least 100 ppm of oleuropein.

11. Olive residue oil which contains at least 300 ppm, preferably at least 350 ppm, more preferably at least 450 ppm of oleanolic acid.

12. Olive oil according to claim 11 which additionally contains at least 20 ppm, preferably at least 100 ppm of oleuropein.

13. Food product, which contains one or more of the following: an olive oil according to claim 9 or 10, an olive residue

oil according to claim 11 or 12, a concentrate obtained according to the process of claim 8.

14. Food product according to claim 13, which is a vegetable oil, preferably an olive oil.

5 15. Food product according to claim 13, which is chosen from the group consisting of spread, mayonnaise, salad dressing or tomato sauce.

10

15

20

25

30

35


40

45

50

55

Figure 1



Oleanolic Acid



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 99 20 4127

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
D,X	L. DI GIOVACCHINO ET AL.: "Effect of mixing leaves with olives on organoleptic quality of oil obtained by centrifugation" JOURNAL OF THE AMERICAN OIL CHEMISTS' SOCIETY., vol. 73, no. 3, 1996, pages 371-374, XP002104605 CHAMPAIGN US * the whole document *	1,3	C11B1/00 A23D9/00 A23D7/00 A23L1/24
A	I. SUSNIK-RYBARSKI ET AL.: "Antioksidaciona svojstva supstancije izolirane iz maslinovog lisca" HRANA I ISHRANA - FOOD AND NUTRITION., vol. 24, no. 1-2, 1983, pages 11-15, XP002134134 BEOGRAD., YU ISSN: 0018-6872 * page 15, Summary *	1,8,13	
A	A. VASQUEZ RONCERO ET AL.: "Acidos triterpenicos del olivo" GRASAS Y ACEITES, vol. 20, no. 3, 1969, pages 133-138, XP002134135 SEVILLA, ES * page 133, column 2, paragraph 2 *	1	TECHNICAL FIELDS SEARCHED (Int.Cl.7)  C11B A23D A23L
A	CA 2 232 692 A (UNIV GRANADA) 5 February 1998 (1998-02-05) * claim 1 *	1,8	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 March 2000	Examiner Dekeirel, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 (P04C01)



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 99 20 4127

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
D,A	<p>DATABASE WPI Section Ch, Week 9722 Derwent Publications Ltd., London, GB; Class B07, AN. 97-241981 XP002104606 - &amp; JP 09 078061 A (TANABE SEIYAKU CO), 25 March 1997 (1997-03-25) * abstract *</p>	1, 8, 13, 14	
A	<p>PATENT ABSTRACTS OF JAPAN vol. 013, no. 227 (C-600), 25 May 1989 (1989-05-25) - &amp; JP 01 039973 A (TAIYO KAGAKU CO LTD), 10 February 1989 (1989-02-10) * abstract *</p>	13	
A	<p>WO 97 32947 A (COMITER TRADING &amp; SERVICES S R ;BALDI ALESSANDRO (IT); ROMANI ANNA) 12 September 1997 (1997-09-12) * claims 1,10,12-16,20 *</p>	9,13,14	
A	<p>US 5 714 150 A (NACHMAN LESLIE) 3 February 1998 (1998-02-03) * claim 1 *</p>	8	
A	<p>A. BIANCO ET AL.: "Microcomponents of olive oil - III: Glucosides of 2(3,4-dihydroxy-phenyl)ethanol" FOOD CHEMISTRY., vol. 63, no. 4, 1998, pages 461-464, XP002134136 ELSEVIER SCIENCE PUBLISHERS LTD., GB ISSN: 0308-8146 * the whole document *</p>	1	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>28 March 2000</b>	Examiner <b>Dekeirel, M</b>
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P4/C01)



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 20 4127

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on .  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

28-03-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
CA 2232692 A	05-02-1998	ES 2111498 A EP 0894517 A WO 9804331 A JP 11513042 T	01-03-1998 03-02-1999 05-02-1998 09-11-1999
JP 9078061 A	25-03-1997	NONE	
JP 01039973 A	10-02-1989	NONE	
WO 9732947 A	12-09-1997	IT F1960041 A EP 0896612 A	05-09-1997 17-02-1999
US 5714150 A	03-02-1998	WO 9938383 A AU 6045098 A	05-08-1999 16-08-1999

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82



The first part of the report deals with the general situation of the country. It is a very interesting and informative study of the country's development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country.

The second part of the report deals with the economic situation of the country. It is a very interesting and informative study of the country's economic development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's economic development.

The third part of the report deals with the social situation of the country. It is a very interesting and informative study of the country's social development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's social development.

The fourth part of the report deals with the political situation of the country. It is a very interesting and informative study of the country's political development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's political development.